Supplement of Earth Syst. Sci. Data, 16, 5681–5701, 2024 https://doi.org/10.5194/essd-16-5681-2024-supplement © Author(s) 2024. CC BY 4.0 License.





## Supplement of

## Microbial plankton occurrence database in the North American Arctic region: synthesis of recent diversity of potentially toxic and/or harmful algae

Nicolas Schiffrine et al.

Correspondence to: Nicolas Schiffrine (nicolas\_schiffrine@uqar.ca)

The copyright of individual parts of the supplement might differ from the article licence.

## Supplementary figures




Figure S1: Number of data samples collected for each Large Marine Ecosystem (LME) region. LME regions are labeled as follows: AI (Aleutian Islands), EBS (East Bering Sea), NB-CS (Northern Bering—Chukchi Seas), BS (Beaufort Sea), CA (Central Arctic), CHA-NG (Canadian High Arctic—North Greenland), CEA-WG (Canadian Eastern Arctic—West Greenland), HBC (Hudson Bay Complex) and L-N (Labrador—Newfoundland).






Figure S2: Number of *Gyrodinium* and *Tripos* occurrence within each Large Marine Ecosystem (LME) region. LME regions are labeled as follows: AI (Aleutian Islands), EBS (East Bering Sea), NB-CS (Northern Bering—Chukchi Seas), BS (Beaufort Sea), CA (Central Arctic), CHA-NG (Canadian High Arctic—North Greenland), CEA-WG (Canadian Eastern Arctic—West Greenland), HBC (Hudson Bay Complex) and L-N (Labrador—Newfoundland).




Figure S3: Proportional contribution of (A) data sources (i.e., SourceArchive; see Table 1) and (B) types of records (i.e., basisOfRecords; see Table 1) across Large Marine Ecosystem (LME) region. The total number of records is given in parentheses next to each region abbreviation. LME regions are labeled as follows: AI (Aleutian Islands); EBS (East Bering Sea); NB-CS (Northern Bering—Chukchi Seas); BS (Beaufort Sea); CA (Central Arctic); CHA-NG (Canadian High Arctic—North Greenland); CEA-WG (Canadian Eastern Arctic—West Greenland); HBC (Hudson Bay Complex); L-N (Labrador—Newfoundland).




Figure S4: Temporal variation in sampling coverage (%) for each Large Marine Ecosystem (LME) region. LME region abbreviations are in parentheses.





Figure S5: Histogram of the distribution of distances to land (in kilometers) for stations sampled in the Beaufort Sea (BS) region. The red dashed line represents the median.



43 Figure S6: Cumulative occurrences (%) of HA taxa.




Figure S7: Geographic distribution of the five most common HA species