

A solar optical hyperspectral library of rare earth-bearing minerals, rare earth oxide powders, copper-bearing minerals and Apliki mine surface samples

Friederike Koerting¹, Nicole Koellner¹, Agnieszka Kuras¹, Nina K. Boesche¹, Christian Rogass¹,
Christian Mielke¹, Kirsten Elger¹, Uwe Altenberger²

¹ GFZ German Research Centre for Geosciences, Potsdam, 14473, Germany
² University of Potsdam, Institute of Geosciences, Potsdam, 14476, Germany

Correspondence to: Friederike Koerting (koerting@gfz-potsdam.de)

Abstract. Mineral resource exploration and mining is an essential part of today’s high-tech industry. Elements such as rare earth elements (REE) and copper are, therefore, in high demand. Modern exploration techniques from multiple platforms (e.g. space- and airborne), to detect and map the spectral characteristics of the materials of interest, require spectral libraries as an essential reference. They include field and laboratory spectral information in combination with geochemical analyses for validation. Here, we present a collection of REE- and copper-related hyperspectral spectra with associated geochemical information. The libraries contain **reflectance** spectra from rare earth element oxides, REE-bearing minerals, copper-bearing minerals and mine surface samples from the Apliki copper-gold-pyrite-mine in the Republic of Cyprus. The samples were measured with the HySpex imaging spectrometers in the visible near infrared (VNIR) and short wave infrared (SWIR) range (400 – 2500 nm). The geochemical validation of each sample is provided with the **reflectance** spectra. The spectral libraries **are** openly available to assist future mineral mapping campaigns and laboratory spectroscopic analyses. The spectral libraries and corresponding geochemistry are published via GFZ Data Services with the following DOIs: <http://doi.org/10.5880/GFZ.1.4.2019.004> (13 REE-bearing minerals and 16 oxide powders, Koerting et al. (2019a)), <http://doi.org/10.5880/GFZ.1.4.2019.003> (20 Copper-bearing minerals, Koellner et al. (2019)), and <http://doi.org/10.5880/GFZ.1.4.2019.005> (37 Copper-bearing surface material **samples** from the Apliki copper-gold-pyrite mine in Cyprus, Koerting et al. (2019b)). All spectral libraries are united and comparable by the internally consistent method of hyperspectral data acquisition in the laboratory.

Deleted: y is

Deleted: (

Deleted: ,

Deleted: .

Deleted: (

Deleted: ,

Deleted: (

Deleted: ,

Deleted: ¶

45

50

60

65

Deleted: study

Deleted: and

Deleted:

Deleted: validation of the materials.

Deleted: for example, for

Deleted: results

Deleted: images

Deleted: serve

Deleted: for the

Deleted: ion

Deleted: ping of

Deleted:

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt


Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

85 Hyperspectral data of geological surfaces can be acquired by ground- or UAV-based outcrop scans to map an ore body's surface mineral distribution by using spectral references libraries. An example of a hyperspectral surface mapping is shown in Figure 1. Here, the outcrop of former copper-gold-pyrite mine Apliki in the Republic of Cyprus was scanned hyperspectrally and mapped utilizing a spectral library of expected surface minerals. The analysis is based on United States Geological Survey (USGS) reflectance spectra. As the USGS spectral library entries do not origin from the same sensor as the mine face scan (HySpex data), they need to be spectrally adapted to the HySpex sensor properties.

90



95 Figure 1: Example for the application of a spectral library. A) 3D model of the open pit Apliki mine in the Republic of Cyprus based on RGB images and a superimposed analysis result of a hyperspectral HySpex scan. The hyperspectral map of the spatial mineral distribution from B) is stacked on the 3D model for visualization purposes. B) Analysis of a HySpex scan using a custom-made spectral library from USGS spectra (Clark et al., 2007). C) Example of hyperspectral spectra from copper-bearing minerals as presented in (Koellner et al., 2019).

Moved (insertion) [1]

Deleted: study

Deleted: surface geometry

Deleted: and

Deleted: (

Deleted:)

Deleted: and can be applied in the laboratory on thin sections, hand samples, as well as drill core to analyse and visualize zonation at a smaller scale.

Deleted: Examples for applications are large-area satellite or aerial surface mapping for geological exploration in early-stage field prospecting. Future hyperspectral imaging satellites will provide the necessary data quality requirements to successfully map rare earth elements (REEs), copper deposits and other resources from space. These satellites will play an important role in the future of geological exploration, to help mapping large mineralized areas in remote regions (Mielke et al., 2016; Swayze et al., 2014). Several global mapping satellite missions will be launched in the next few years. Amongst them are the German EnMAP, the Chinese CCRSS-A and the Japanese HISUI missions (Guanter et al., 2015; Iwasaki et al., 2011; Tong et al., 2014). For those missions, the imaging spectroscopy community is currently developing methodologies for e.g. the detection of REEs in the image spectra (Boesche et al., 2015; Boesche, 2015; Bösche, 2015; Herrmann, 2019; van der Meer et al., 2012; Turner et al., 2014a, 2014b; Turner, 2015). Hyperspectral data can be acquired by ground- or UAV-based outcrop scans to study an ore body's surface geometry and mineral distribution (Figure 1) and can be applied in the laboratory on thin sections, hand samples, as well as drill core to analyse and visualize zonation at a smaller scale.

Moved up [1]: Hyperspectral data can be acquired by ground- or UAV-based outcrop scans to study an ore body's surface geometry and mineral distribution (Figure 1) and can be applied in the laboratory on thin sections, hand samples, as well as drill core to analyse and visualize zonation at a smaller scale.

Deleted: ling

Deleted: 79

Deleted: one

Deleted: RGB

Deleted: ene

Deleted: from the Apliki mine in the Republic of Cyprus.

Deleted: mineral

Deleted: analysis

Deleted:

Deleted: based on

Deleted: ene

Deleted: analyzed by EnGeoMap 2.0

Deleted:

Deleted: (Mielke et al., 2016)

Formatted

145 In the case of minerals reflectance spectra, only hyperspectral sensors with a spectral bandwidth resolution of approximately 10nm or less can capture the fine differences in reflectance at certain wavelength positions (Jensen, 2010). Future hyperspectral imaging satellites will provide the necessary data quality requirements to successfully map rare earth elements (REEs), copper deposits and other resources from space. These satellites will play an important role in the future of geological exploration, to help mapping large mineralized areas in remote regions (Mielke et al., 2016; Swayze et al., 2014).

150 Several global mapping satellite missions will be launched in the next few years. Amongst them are the German EnMAP, the Chinese CCRSS-A and the Japanese HISUI missions (Guanter et al., 2015; Iwasaki et al., 2011; Tong et al., 2014). For those missions, the imaging spectroscopy community is currently developing methodologies e.g. for the detection of REEs in the image spectra (Boesche et al., 2015; Boesche, 2015; Bösche, 2015; Herrmann, 2019; van der Meer et al., 2012; Turner et al., 2014a, 2014b; Turner, 2015).

155 We aim to contribute to the already existing, accredited libraries, e.g. the USGS- and the ECOSTRESS Spectral Library and various others (Baldrige et al., 2009; Clark et al., 2007; Hunt, 1977; Kokaly et al., 2017; Meerdink et al., 2019; Percival et al., 2016). The available reflectance spectral libraries are commonly based on powdered natural or synthetic samples that are spectrally pure. The spectral data is usually collected by point-spectroradiometers e.g. the Analytical Spectral Device (ASD) FieldSpec® 3. Our contributed reflectance spectra are based on imaging spectroscopy data from the HySpex classic series scanning samples in a natural and a powdered state. Reflectance spectral libraries like the here presented, based on HySpex imaging data and untreated samples, are not yet freely available for the hyperspectral community.

160 The spectral and geochemical information of samples presented here, belong to three different mineral assemblages and correspond to three different types of deposits. The sample's spectral information is provided within four spectral library files and their corresponding geochemical composition files. The four spectral library files represent (1) REE-bearing minerals, (2) synthetic REE-oxide powders (Koerting et al., 2019a), (3) copper-bearing minerals (Koellner et al., 2019) and (4) powders of copper-bearing surface material from the Apliki copper-gold-pyrite mine in the Republic of Cyprus (Koerting et al., 2019b). Spectrally, the libraries cover the full wavelength range of the solar optical range (414 nm – 2498 nm). The corresponding geochemical analyses are explained in the methods for each sample type.

170 The two REE libraries (Koerting et al., 2019a) consist of the spectra of 16 rare earth oxide (REO) powders and 13 REE-bearing minerals (REMin). In addition, the spectra of niobium- and tantalum oxide powders are provided, which will further not be mentioned individually but be included in the term "REO". The third spectral library includes 20 copper-bearing minerals (Koellner et al., 2019) and the fourth spectral library contains 37 surface samples from the Apliki copper-gold-pyrite mine site in the Republic of Cyprus (Koerting et al., 2019b). All spectral libraries are united and comparable by the internally consistent method of hyperspectral data acquisition in the laboratory. An extensive list of the samples can be found in the technical reports provided with each dataset.

175

Deleted: The methods developed for the upcoming hyperspectral satellite missions like EnMap (Guanter et al., 2015) use image spectra as a database for their analysis.

Formatted: Font: Font colour: Black, German

Formatted: German

Deleted: The image scenes are acquired by a moving line scanner mounted on the satellite, which records the spatial dimension (x- and y-dimension) line by line, as well as the wavelength dimension (z-dimension). Each pixel therefore represents the full spectral range of the sensor. The sensor's movement along a rotation or a movement axis provides spatially continuous imaging spectroscopy data. Variations along the spectral domain of the data are visible as concave indentions, often referred to as "absorption bands". They are characteristic for the measured surface material and enable its identification and quantification (Clark, 2003). The recorded spectral information is a function of the chemical and physical properties of the target material and the sensor setup itself. Typical copper-bearing hyperspectral spectra are shown in Figure 1C.

We are presenting four spectral library files and their geochemical validation. The library files present the spectral information of three different mineral groups: (1) REE-bearing minerals and (2) synthetic REE-oxide powders (Koerting et al., 2019a), (3) copper-bearing minerals without any sample preparation (Koellner et al., 2019) and (4) powders of copper-bearing surface material from the Apliki copper-gold-pyrite mine in the Republic of Cyprus with the corresponding GPS position of the samples (Koerting et al., 2019b). Spectrally, the libraries cover the full wavelength range of the solar optical range (414 nm – 2498 nm). The geochemical validation for each sample type is explained in the methods....

Moved (insertion) [2]

Deleted:

Moved up [2]: The geochemical validation for each sample type is explained in the methods.

Deleted: ds.

Deleted: 4

Deleted:

Deleted:

Deleted: all

Deleted: supplements

215 The samples are presented as reflectance spectral libraries and their geochemical composition. Sample nominations are based on the geological collection of origin or sample abbreviations from the field sampling. The sample nomination is not an interpretation of the presented geochemical data. The datasets are independent of each other, the reflectance spectra can be seen as a spectral expression of the existing geochemical data. Neither the geochemistry nor the reflectance spectra are interpreted or correlated to each other.

220 The outline of this document follows the necessary line of knowledge to successfully make use of the here presented spectral libraries. Section 2 includes a description of the analysed materials and Section 3 informs about the methods including the sample preparation and spectra collection, the hyperspectral data acquisition, covering the processing of the data and spectral measurement parameters and the geochemical analyses of the samples. Section 4 lists the samples that were measured spectrally and geochemically and the data of which can be accessed via the GFZ Data Services platform. Section 5 discusses the parameters influencing the data. A separate data description and the geochemical analysis results are included as data reports in the three different data publications (Koellner et al., 2019; Koerting et al., 2019a, 2019b).

2. Materials

230 The REE sample material includes 16 REO powders (REO) and 13 REE-bearing minerals (REMin). The REO powders belong to a series of rare earth metals and compounds (REacton®) and were purchased from Alfa Aesar. All REO powders contained at least 99.9% of the REE oxide, as per the seller-supplied concentration certificates. The concentration certificate information can be found in the data description of (Koerting et al., 2019a). The REO powders were obtained as high-purity materials with a grainsize of <63 µm. The REMin samples (ore minerals) were purchased from Gunnar Färber Minerals, an online trader of mineral specimens. The mineral notation is based on the sample name provided by Gunnar Färber Minerals.

235 The supplier offers analytical services with a modern REM-EDX technology and therefore we assume the specimen are analysed and the mineral species is validated before the sale. The X-Ray Fluorescence (XRF) data presented in the data description of (Koerting et al., 2019a) should be consulted to validate the given mineral nomination noted by Gunnar Färber Minerals.

240 The 20 copper-bearing minerals belong to collections of the University of Potsdam (UP) and the Federal Institute for Geosciences and Natural Resources (BGR), a samples list can be found in (Koellner et al., 2019). The minerals were measured hyperspectrally with no sample preparation, the sample photos and geochemical analysis are provided in the data description for (Koellner et al., 2019). The 37 Apliki mine surface samples were collected (Koerting et al., 2019b) in March 2018 during a field campaign of the Geological Survey Department of the Republic of Cyprus (GSD) and the GFZ German Research Centre for Geosciences (GFZ). Surface material in the mine was collected and prepared (crushed and pulverized) for the geochemical analysis by Bureau Veritas Minerals (BVM). The powdered samples were measured hyperspectrally as

Deleted:

Deleted: z

Deleted: . Section 4 describes

Deleted: Section 5 presents

Deleted: Section 6 discusses the parameters influencing the data.

Deleted: The spectral libraries can be downloaded as an ASCII *.txt file format and as binary spectral library files format *.sli with its associated header files *.hdr .

Deleted: .

Commented [FK1]: Add table listing all sample names, IDs and mineral names if possible

Deleted: ¶

Deleted: 4

Deleted: his

Deleted: z

Deleted: A sample list for the REMin can be found in .S1 and S2, for the REO in S3 of the Supplement.¶

Formatted: Justified, Space After: 0 pt, Widow/Orphan control

Deleted: the appendix (S4-S5).

powder tablets, a sample list including photos from the in-situ conditions of the samples can be found in [the technical report](#) (Koerting et al., 2019b).

3. Methods

3.1 Sample Preparation and spectra collection

The sample preparation varies by sample type and depends on the material and the information of interest. This is based on the research projects that the samples stem from and for which the spectral and geochemical data was acquired.

The reflectance spectra for each sample were manually extracted from the processed hyperspectral image scenes by averaging a number of pixels over a central sample area. The resulting spectra were compiled in a spectral library. Thereby, each reflectance spectrum of a spectral library represents an average reflectance spectrum of the material, depending on the sample size and spectral homogeneity. The extraction of the reflectance spectra is explained in detail in each data description (Koellner et al., 2019; Koerting et al., 2019a, 2019b).

The REO powders were measured in 100% quartz glass petri dishes underlain by black cellular rubber, each powder was measured separately. Figure 2 shows the measurement setup of holmium-oxide powder as an example for the REO powders. The REE-bearing minerals were measured separately. Figure 3 shows the xenotime sample (brownish single crystal embedded in quartz) as an example for the REMin samples. The REMin samples were measured without sample preparation on black cellular rubber, as is shown for the copper-bearing minerals in Figure 4. For all measurements, the final reflectance spectral analyses were spatially reduced to the centre pixels of each identified REE-bearing mineral or a 5x5 pixel average reflectance spectrum centred on the REO powder sample. Shadow effects from the sidewalls of the boxes could thus be minimized. One representative reflectance spectrum of every REMin and REO sample was collected for the spectral library (Herrmann, 2019).




Figure 2: Holmium-oxide powder in the laboratory HySpex setting in a quartz glass petri dish underlain by black cellular rubber. Geometric markers for the pre-processing were placed alongside the sample.

Deleted: S6 of the supplement and

Deleted: in

Deleted: d

Deleted: s

Deleted: using an

Deleted: and

Deleted: .

Deleted: averaging

Deleted: erefore

Deleted: produced

Deleted: . In order to improve the signal-to-noise ratio and reduce the influence of outlier pixels, an average spectrum of sample-covering pixels was produced ...



Figure 3: Xenotime embedded in quartz as an example for the REE-bearing mineral samples.

The copper-bearing mineral samples were measured without any sample preparation as the variable surface of the minerals and the influence of the mineral structure was of interest. Figure 4 shows an example scan of some of the copper-bearing minerals. The full sample list including sample photos and the marked area of the geochemical sampling can be found in the technical report (Koellner et al., 2019). The area, used to obtain the spectrum, averaging over a 5x5 pixel window, was afterwards sampled for the geochemical analysis.



Figure 4: Showing HySpex scan “MH_FK_LAB_Cudetect_008_09012018_WR20” as an example to highlight the lack of sample preparation.

The Apliki mine samples were crushed and powdered so that ≥85% of the sample was below 75µm. Homogenized powders were measured as pressed powder tablets (Figure 5). The area to obtain the sample’s reflectance spectrum was chosen over a 5x5 pixel window in the centre of the powder tablet to minimize influences from the tablet’s metal frame. The dark spots in each tablet were caused by previous measurements with a laser induced breakdown spectrometer (LIBS). The hyperspectral

Deleted: ¶
Deleted: no
Deleted: of a part
Deleted: Table S5 in the supplements shows the area for the geochemical sampling encircled in yellow. The same
Deleted: same
Deleted: was
Deleted: by
Deleted:
Deleted: .

Deleted: <
Deleted: a
Deleted: by
Deleted: ,
Deleted: .

sample spots were chosen in order to exclude the measurement points of the LIBS in the spectral footprint. In case of broken powder tablets like “7d_Hem”, the shadowed, rough surface areas were excluded from the spectral sampling and an even powder surface was favoured.




Figure 5: Showing the Apliki mine samples prepared as powder tablets.

3.2 HySpex Data Recording

The HySpex VNIR-1600 and SWIR-320m-e (technical description available at: (hyspex.no/products/disc.php, 2019)) are two line-scanning cameras mounted in parallel. They cover the range of the visible to near infrared (VNIR, 414 – 993 nm) and the short-wave infrared (SWIR, 967 – 2498 nm) wavelength region. The sensors record an array-line of 1600 pixel (VNIR) and 320 pixel (SWIR) (push-broom scanning). Every pixel contains a spectrum with a total spectral sampling number of 408 bands in total.

The HySpex cameras are provided with two acquisition modes, one for airborne data collection and one for laboratory measurements. In laboratory mode, the cameras are combined with a trigger pulse-moving sleigh (translation stage) of definable frame period (depending on the integration time of every array-line acquisition). The configuration of the translation stage framework, the cameras and the light source (Halogen GX6.35, 2 x 1000 W, 45° illumination angle) are fixed, while the sleigh and the samples are moving through the focal plane (Rogass et al., 2017).

The reflectance level of a white reference panel, placed in line with the samples, is chosen according to the albedo of the samples. The higher the albedo of the sample, the higher is the diffuse reflectance factor of the white reference panel that is chosen. For the REE samples (REMin and REO), a white reference panel of 95% reflectance was used, because most of the REO samples were bright, white powders of a high albedo, this is based on test measurements of (Bösche, 2015; Herrmann, 2019). The Apliki samples required a 50% reflectance white reference panel, whereas the copper-bearing minerals were measured using a 20% reflectance white reference panel. Both, the geometrical setup and the heat up time of the lamp influence the configuration of the light source. The maximum illumination was obtained with an angle of 45° between the

Deleted: also

Deleted: .

Deleted: 4.

Deleted: 1000

Deleted: 1000

Deleted: y

Deleted: certai

370

375

380

incident light and the vertical plane. The distance between the lamp and the HySpex cameras was higher compared to the distance between the samples and the sensor to ensure diffuse illumination and to avoid thermal influence on the cameras and the samples. The integration time (= measurement time for each image line) was tested to be as high as possible to suppress the impact of signal uncorrelated gaussian white noise and at the same time as low as needed to avoid detector saturation. For all measurements the integration time was chosen with respect to the sample albedo. The HySpex sensor characteristics are listed in Table 1. The used settings for the REMin and REOs are listed in Table 2, the settings for the copper-bearing minerals in Table 3, and for the Apliki mine samples in Table 4. The laboratory is equipped with black-painted walls and doors, as well as black curtains to avoid reflected light from surfaces other than the sample, an example setup of the sensors, the translation stage and the samples can be seen in Figure 6. The laboratory conditions were kept stable, the air temperature was regulated to $21\pm0.5^{\circ}\text{C}$ and the humidity was below 70% for all measurements. Black cellular rubber is used as a base material for all samples for hyperspectral data acquisition. It reflects less than 5% on average of the incoming radiation.

Detailed descriptions for the GFZ' standard measurements and the process chain can be found in (Rogass et al., 2017).




Figure 6: The HySpex translation stage setup (Körting, 2019).

385

Deleted: 2

Deleted: 3

Deleted: 4

Formatted: Caption, Space After: 0 pt, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: Not at 9,1 cm

Deleted: ¶

390

Table 1: HySpex sensor parameters of the VNIR-1600 (VNIR) and SWIR-320m-e (SWIR)

HySpex sensor parameters		
Lamp arrangement	45°	
	VNIR	SWIR
Wavelength range [nm]	414 - 993	967 - 2498
Pixels per line	1600	320
Sampling interval [nm]	3.7	6
Radiometric resolution	12 bit	14 bit
Light source	Halogen GX6.35, 2 x 1000 W	

Table 2: HySpex settings for laboratory measurements of the REO and REMin (Koerting et al., 2019a, modified after (Bösche, 2015; Herrmann, 2019).

HySpex settings		
Distance, sample to sensor	1 m	
Sensor arrangement head to head	1m lenses, eq on VNIR	
	VNIR (1600 px)	SWIR (320 px)
Integration time [µs]	30 000	5 000
Frame period [µs]	31 000	123 506

Table 3: HySpex settings for laboratory measurements of the copper-bearing minerals (Koellner et al., 2019).

HySpex settings		
Distance, sample to sensor	30cm	
Sensor arrangement head to head	30cm lenses, eq on VNIR	
	VNIR (1600 px)	SWIR (320 px)
Integration time [µs]	120000 - 140000	15000 – 20000
Frame period [µs]	120062 - 141004	478334 - 561768

Table 4: HySpex settings for laboratory measurements of Apliki mine powdered samples (Koerting et al., 2019b).

HySpex settings		
Distance, sample to sensor	1 m	
Sensor arrangement head to head	1m lenses, eq on VNIR	
	VNIR (1600 px)	SWIR (320 px)
Integration time [µs]	60000	10000
Frame period [µs]	60060	239282

395

Formatted: Tab stops: 4,68 cm, Left

Formatted Table

Deleted: 2

Deleted: Lamp arrangement ... [1]

Deleted: Wavelength range ... [2]

Deleted: Frames ... [3]

Deleted: ¶

Deleted: 3

Deleted: Lamp arrangement ... [4]

Deleted: Wavelength range ... [5]

Deleted: Frames ... [6]

Deleted: ¶

Deleted: 4

Deleted: Lamp arrangement ... [7]

Deleted: Wavelength range ... [8]

Deleted: Frames ... [9]

Deleted: ¶

420 **3.3 Hyperspectral Data Processing**

Each measurement run produces one VNIR and one SWIR 3D-data cube. The three dimensions are the two spatial x,y- and the spectral z-dimension. The 3D image cubes are produced, by moving a homogeneous reflecting white reference panel and the samples through the focal plane of the two sensors. The VNIR image cube is resized to the spatial dimensions of the SWIR data cube, co-registered and stacked with the SWIR data cube resulting in a continuous image cube with the spectral range of 414 – 2498 nm. In order to produce a reflectance image, the image pixel that show the white standard were averaged to a one-line reference spectrum. The reflectance was calculated by dividing every image line spectrum by its reference spectrum from the reflecting white reference panel. A detailed description for the laboratory set-up and processing can be found in (Rogass et al., 2017). The software ‘HySpex ground’ was used to perform the measurements and the software ‘HySpex rad’ was used to perform the radiometric calibration on the image data.

430 **3.4 Geochemical Sample Analysis for Sample Characterization**

Depending on the sample type, the geochemical analysis methods differ. The methods used for each sample type, are listed in Table 5.

Table 5: Sample type and corresponding geochemical characterization method.

Sample type	Geochemical Analysis
REO (Koerting et al., 2019a)	Laboratory certificates
REMin (Koerting et al., 2019a)	X-Ray Fluorescence (XRF), Electron probe microanalyzer (EPMA) analyses
Copper-bearing minerals (Koellner et al., 2019)	Scanning electron microscope (SEM), EPMA
Apliki mine samples (Koerting et al., 2019b)	Bureau Veritas Mineral Analysis, ICP-MS and ES

435 **3.4.1 Thermo Niton XL3t (XRF)**

REMin

The geochemical measurements for the REMins were performed using an X-Ray Fluorescence (XRF) instrument - Thermo Niton XL3t (Fisher Scientific, 2002). The XL3t is a lightweight, hand-held XRF analyzer. The measurement principle follows the principle of X-Ray fluorescence, where the sample inbound X-Rays excite electrons to a higher energy level in the sample material. Energy in form of XRF radiation is released when these electrons return to their original state. The frequency of this radiation is characteristic for the measured chemical element and its intensity is correlated to the concentration level. The intensity of each element is detected as counts per second by the detector, a geometrically optimized

large area drift detector (GOLDD). The maximum excitation voltage of the XL3t device is 50 kV, which means out of the full REE suite only four light REEs can be detected (Lanthanum, Cerium, Praseodymium and Neodymium).

The XL3t spectrometer is attached to a lead shielded sample chamber, in which samples with a diameter smaller than 3.3 cm can be placed. Mineral samples can be directly placed in the chamber; powdered samples have to be placed in sample tubes (2.5 cm diameter). The sample tubes are made of plastic with a plastic foil on the bottom. The plastic cannot be detected by XRF and therefore not interfere with the measurements. A built-in camera of the XL3t enables the precise location of the measuring spot. The software used for the measurements is named “NDTr” and the measurement mode was “mining and exploration”. The concentration levels are provided along with a balance value. “Balance” represents counts per seconds that could not be attributed to one of the measured elements. Table 6 shows the measurement modes and filters used. In-depth description of the XL3t and the XL3t-results for each sample can be found in (Bösche, 2015; Herrmann, 2019)

Deleted:

Deleted: 6

Table 6: Settings used for the Thermo Niton XL3t X-ray fluorescence device (Bösche, 2015).

Deleted: 6

Thermo Niton XL3t Setting	
Measurement mode	Test all geo
Filter	Main, Low, High, Light
Filter measurement time	30 seconds each

3.4.2 Scanning Electron Microscope (SEM) and Electron Probe Microanalyzer (EPMA)

Deleted: 5.2

Copper-bearing minerals

In order to obtain information about the zonation and internal fabrics of the copper-bearing minerals a fully automated JEOL JSM-6510 scanning electron microscope (SEM) (20kV acceleration voltage) at the University of Potsdam was used. A back-scattered electron detector displays compositional variation in the imaging area based on the mean atomic number of the pixel. An energy dispersive X-ray spectrometer (EDX, Oxford Instruments INCAX-act) attached to the instrumentation provides quantitative elemental analysis of single spots. After calibrating with pure copper, a wide spectrum of elements can be identified. Based on previous results, divergences of up to 5 weight % can be expected, which for quantitative analysis is acceptable.

In order to approximate the values for copper a JEOL JXA-8200 electron probe microanalyzer (EPMA) at the University of Potsdam was used. The electron microprobe is equipped with five wavelength-dispersive X-ray spectrometers (WDX) and was operated with a 20 kV accelerating voltage, a 20 nA current, and a beam diameter of 2 µm. The analytical counting times were 20/10 s for the element peak and 10/5 s for background positions. Analyses were calibrated using silicates/sulphides obtained from the Smithsonian Institution and Astimex. Quantifying elements of a lower atomic mass than boron is not possible, carbon cannot be measured either.

An example SEM analysis for copper-bearing mineral sample “C1 Chalcopyrite” can be seen in Figure 7, the EPMA analysis of the mineral is listed in Table 7. The full SEM and EPMA results are documented in (Koellner et al., 2019).




Figure 7: Sample C1 Chalcopyrite SEM and EPMA analysis.

Table 7: Sample C1 Chalcopyrite EPMA analysis results, from three sample-points on the sample. Element concentrations reported in wt% or as “below detection limit” (bdl).

Sample-point	Al [wt%]	Hg [wt%]	Fe [wt%]	Cu [wt%]	Si [wt%]	S [wt%]	Mn [wt%]	Total [wt%]
C1_Chalcopyrite-1	bdl	bdl	30.00	33.98	bdl	34.81	bdl	98.79
C1_Chalcopyrite-2	bdl	bdl	30.19	34.108	bdl	34.94	bdl	99.23
C1_Chalcopyrite-3	bdl	bdl	30.08	34.194	bdl	35.09	bdl	99.36

REE-bearing Minerals

Some of the REMin (xenotime, bastnaesite, fluorapatite, synchysite and ilmenite) were additionally analysed by using a JEOL JXA-8200 electron microprobe (EPMA) at the University of Potsdam based on a method developed by (Lorenz et al., 2019). The conditions used for the analysis were: 20kV acceleration voltage, 20nA beam current and a beam size of 2 µm. Counting times were between 10 s - 20 s on peak for major elements and 50 s for REE and other trace elements.

The following spectral lines and mineral standards from Smithsonian and Astimex were used: fluorapatite (F K α , P K α , Ca K α), albite (Na K α), fayalite (Fe K α , Mn K α), wollastonite (Si K α), omphacite (Al K α), LaPO $_4$ (La L α), PrPO $_4$ (Pr L β), CePO $_4$ (Ce L α), NdPO $_4$ (Nd L β), YPO $_4$ (Y L α), EuPO $_4$ (Eu L α), SmPO $_4$ (Sm L β), LuPO $_4$ (Lu L α), GdPO $_4$ (Gd L α), ErPO $_4$ (Er L β), DyPO $_4$ (Dy L β), YbPO $_4$ (Yb L α), HoPO $_4$ (Ho L β), uranotorite (U M β), crocoite (Pb M β). The EPMA data were

Deleted: and Table 7,

Deleted: t

Deleted: data files

No: 16

backscatter electron image

C1 - S55L16C (Mount 1)

Deleted:

Formatted: Normal

Deleted: example

Deleted: ¶

Formatted: Caption, Tab stops: Not at 9,98 cm

Deleted: 7

Deleted:

Deleted: example

Deleted: up to

Deleted: 3

Deleted: measurement

Deleted: spots for each

Deleted: (C1-S55L16c-1 – 3 in the “comment column”).

Deleted: as

Deleted: .

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Deleted: → ¶

Al ... [10]

Deleted: z

Deleted: monazite (Th Ma, U Mβ, Tb La),

reduced using the software-implemented PRZ-XXP data-correction routine, which is based on the $\phi(\rho z)$ method (Heinrich and Newbury, 1991).

530 **3.4.3. Apliki mine surface sample analysis**

The Apliki mine samples were analysed by Bureau Veritas Minerals – Mineral Laboratories Canada (BVM) using their standard packages (Bureau Veritas, 2020). The samples were pulverized below 75 μm and analysed for major, minor and trace elements using ICP-MS and ES. The results are grouped by the internal BVMs sample preparation-/ analysis method types. Those “analysis method types” were namely “aquatic”, “rock” and “soil”. The sample numbers, associated analysis method, type and internal BVM analysis codes can be found in the technical report of the Apliki mine surface sample data (Koerting et al., 2019b).

540 **4. Results**

The following samples are provided and described in detail in the corresponding technical reports. For clarity purposes, all provided samples and corresponding spectra names are listed here, including a short sample description and, where applicable, the sampling location, geochemistry or mineralogy (Table 8-12). The detailed sample descriptions can be found in the corresponding technical reports provided with the data. For each file collection a plot of the spectral library is shown (Figure 8-13).

545 **4.1 REE-bearing minerals and rare earth oxide powders**

Table 8: Samples, sample names and locality and spectral library filenames of REE-bearing minerals

Sample	Original sample name	Sample locality	Spectrum name
Aeg	Aegirine, "Acmite"	Rundemyr, Øvre Eiker, Buskerud, Norway/TYP	REMin_Aeg
Bar	Bariopyro-chlore, Fluorapatite	Mina Boa Vista, Catalao, Goias/Brazil	REMin_Bar
Bst	Bastnaesite (Ce)	Zagi Mountain, Warzal Dam, Pechawar, North-West Frontier Prov./Pakistan	REMin_Bst
Fap	Fluorapatite, Albite	Golconda Mine, Governador Valadares, Doce Valley, Minas Gerais/Brazil	REMin_Fap
Flt	Fluorite	Arbegona, Shashemange	REMin_Flt
Gdl	Gadolinite (Y) Synchysite (Y), Fluorite	White Cloud Pegmatite, South Platte, Jefferson Co. Colorado/USA	REMin_Gdl
Ilm	Ilmenite	Mogok, Sagaing District, Mandalay/Myanmar	REMin_Ilms
Per	Polyrase (Y)	Puoutevare Peg-matite, Tjalmajaure Lake, Jokkmokk Lappland/Northern Sweden	REMin_Per
Prs	Parisite (Nd) incl. Parisite (Ce)	Mountain Pass Mine, Ivanpah Mts. San Bernardino Co. California/USA	REMin_Prs
Syn	Synchysite (Y), Microcline, Quartz	White Cloud Pegmatite, South Platte, Jefferson Co. Colorado/USA	REMin_Syn
Xtm1	Xenotime (Y) (a)	Novo Horizonte, Ibitiara, Bahia/Brazil	REMin_Xtm1
Xtm2	Xenotime (Y) (b)	Novo Horizonte, Ibitiara, Bahia/Brazil	REMin_Xtm2
Zrn	Zircon	Peixe alkaline complex, Monteirópolis, Jaú do Tocantins, Tocantis/Brazil	REMin_Zrn

Deleted: 5.3	
Deleted: zed by Bureau Veritas Minerals – Mineral Laboratories	[11]
Formatted	... [12]
Formatted	... [13]
Formatted	... [14]
Formatted	... [15]
Formatted	... [17]
Formatted Table	... [16]
Formatted	... [18]
Formatted	... [19]
Formatted	... [20]
Formatted	... [21]
Formatted	... [22]
Formatted	... [23]
Formatted	... [24]
Formatted	... [25]
Formatted	... [26]
Formatted	... [27]
Formatted	... [28]
Formatted Table	... [29]
Formatted	... [30]
Formatted	... [31]
Formatted	... [32]
Formatted	... [33]
Formatted	... [34]
Formatted	... [35]
Formatted	... [36]
Formatted	... [37]
Formatted	... [38]
Formatted Table	... [39]
Formatted	... [40]
Formatted	... [41]
Formatted	... [42]
Formatted	... [43]
Formatted	... [44]
Formatted	... [45]
Formatted	... [46]
Formatted	... [47]
Formatted Table	... [48]
Formatted	... [49]

Table 9: Sample name and supplier, product and lot number and spectral library filenames of the rare earth oxide powders.

Sample name, Supplier	Product Number	Lot Number	Spectrum Name
Yttrium (III) oxide, Sigma-Aldrich	204927	MKBL2030V	REO_Yttrium
Niobium (V) oxide, Alfa Aesar	11366	L18Y022	REO_Niobium
Lanthanum (III) oxide, Alfa Aesar	11272	B08X015	REO_Lanthanum
Cerium (IV) oxide, Alfa Aesar	11372	L07S057	REO_Cerium
Neodymium (III) oxide, Alfa Aesar	11250	C02W029	REO_Neodymium
Samarium (III) oxide, Alfa Aesar	11229	61200836	REO_Samarium
Europium (III) oxide, Alfa Aesar	11299	A16Z001	REO_Europium
Gadolinium (III) oxide, Alfa Aesar	11290	A13W016	REO_Gadolinium
Terbium (III,IV) oxide, Alfa Aesar	11208	J24Q019	REO_Terbium
Dysprosium (III) oxide, Alfa Aesar	11319	61300733	REO_Dysprosium
Holmium (III) oxide, Alfa Aesar	11280	J11X030	REO_Holmium
Erbium (III) oxide, Alfa Aesar	11310	61000356	REO_Erbium
Thulium (III) oxide, Alfa Aesar	11198	E25S060	REO_Thulium
Ytterbium (III) oxide, Alfa Aesar	11191	61201069	REO_Ytterbium
Lutetium (III) oxide, Alfa Aesar	11255	G14X082	REO_Lutetium
Tantalum (V) oxide, Alfa Aesar	14709	114Y039	REO_Tantalum




Figure 8: Spectral library plot of the REE-bearing minerals.

Formatted	... [50]
Formatted	... [51]
Formatted	... [53]
Formatted Table	... [52]
Formatted	... [54]
Formatted	... [55]
Formatted	... [56]
Formatted	... [57]
Formatted	... [58]
Formatted	... [59]
Formatted	... [60]
Formatted	... [61]
Formatted	... [62]
Formatted	... [63]
Formatted	... [64]
Formatted	... [65]
Formatted	... [66]
Formatted	... [67]
Formatted	... [68]
Formatted	... [69]
Formatted	... [70]
Formatted	... [71]
Formatted	... [72]
Formatted	... [73]
Formatted	... [74]
Formatted	... [75]
Formatted	... [76]
Formatted	... [77]
Formatted	... [78]
Formatted	... [79]
Formatted	... [80]
Formatted	... [81]
Formatted	... [82]
Formatted	... [83]
Formatted	... [84]
Formatted	... [85]
Formatted	... [86]
Formatted	... [87]
Formatted	... [88]
Formatted	... [89]

575




Figure 9: Spectral library plot of the rare earth oxide powders.

4.2 Copper-bearing minerals

Table 10: Sample names, collection, original sample name, locality, alteration, mineral formula, spectral library filenames and geochemical composition of the copper-bearing sulphides and native copper

Sample name	Collection	Original sample name	Sample locality	Visible alteration	Spectra names	Geochemical composition (EPMA mean, n=3, wt%)
C1 Chalcopyrite	BGR	S55L16 C	Fösseberg Mine, Siegerland, Germany	strongly altered	C1 Chalcopyrite_BG R-S55L16-C [5x5 AVG]	S: 34,941; Fe: 30,091; Cu: 34,094
C2 Chalcopyrite	BGR	S115R12	Erzgebirge, Slovakia	slightly altered	C2 Chalcopyrite_BG R-S115R12 [5x5 AVG]	S: 34,903; Fe: 30,068; Cu: 33,95
C3 Chalcopyrite	BGR	S131L5 C	Henderson Mine, Clear Creek Country, USA	tarnished	C3 Chalcopyrite_BG R-S131L5-C [5x5 AVG]	S: 35,039; Fe: 30,106; Cu: 33,965
C4 Chalcopyrite	UP	7534	Cornwall, England, GB	slightly altered	C4 Chalcopyrite_UP-7534 [5x5 AVG]	S: 35,007; Fe: 30,156; Cu: 34,044
C5 Chalcopyrite	UP	7526	Clausthal, Harz, Germany	altered	C5 Chalcopyrite_UP-7526 [5x5 AVG]	S: 35,053; Fe: 30,007; Cu: 34,177
K1 Copper	UP	600-1	Furnace, Lübeck, Germany	slightly altered	K1 Copper_UP-600-1 [5x5 AVG]	Cu: 98,577

580

Table 11: Sample names, collection, original sample name, locality, alteration, mineral formula, spectral library filenames and geochemical composition of the copper-bearing silicates, carbonates and sulphates.

Sample name	Collection	Original sample name	Sample locality	Visible alteration	Spectra name	Geochemical composition (EPMA mean, n=3, wt%)
A1 Azurite	UP	2458	Cheroy near Lyon, France	altered, nodular	A1 Azurite UP-2458 [5x5 AVG]	CuO: 65.344; HgO: 0.091
A2 Azurite	UP	2437	Tsumeb near Otavi, Namibia	altered	A2 Azurite UP-2437 [5x5 AVG]	CuO: 65.194
A3 Azurite	BGR	S101L7	Cornberg by Fulda, Germany	strongly altered	A3 Azurite BGR-S101L7 [5x5 AVG]	CuO: 63.87; SO ₃ : 0.127; FeO: 0.179
B1 Brochantite	BGR	S115R3	Altenberg, Slovakia	slightly altered, powdered	B1 Brochantite BG-R-S115R3 [5x5 AVG]	Al ₂ O ₃ : 0.18; SiO ₂ : 0.069; SO ₃ : 16.262; CuO: 80.334
F1 Unknown	BGR	S115R14	Kotterbach near Witkow, Poland	slightly altered	F1 Unknown BGR-S115R14 [5x5 AVG]	SiO ₂ : 2.588; FeO: 69.042; CuO: 0.25; SO ₃ : 0.161; MnO: 0.292
L1 Linarite	UP	9542	Unknown location	slightly altered, acicular	L1 Linarite UP-9542 [5x5 AVG]	SO ₃ : 64.18; CuO: 24.184; HgO: 0.439
M1 Malachite	BGR	S134R8	L'Etoile du Congo Mine, Katanga, Kongo	altered, nodular	M1 Malachite BGR-S134R8 [5x5 AVG]	CuO: 67.609
M2 Malachite	BGR	S131L5 M	Henderson Mine, Clear Creek Country, USA	strongly altered	M2 Malachite BGR-S131L5-M [5x5 AVG]	CuO: 66.917
M3 Malachite	BGR	S131R4	Tsumeb near Otavi, Namibia	altered	M3 Malachite BGR-S131R4 [5x5 AVG]	CuO: 65.176; SO ₃ : 0.458
M4 Malachite	BGR	S132L2	Ogonja Mine in Okavandja, Namibia	strongly altered	M4 Malachite BGR-S132L2 [5x5 AVG]	CuO: 67.051
M5 Malachite	BGR	S55L16 M	Siegen, Germany	slightly altered, acicular	M5 Malachite BGR-S55L16-M [5x5 AVG]	CuO: 67.885
P1 Plancheite	UP	Oberhä2	Jordan	slightly altered	P1 Plancheite UP-Oberhä [5x5 AVG]	Al ₂ O ₃ : 2.951; SiO ₂ : 42.079; CuO: 51.782; SO ₃ : 0.061; MnO: 0.243
P2 Plancheite	UP	Oberhä2	Jordan	slightly altered	P2 Plancheite UP-Oberhä2 [5x5 AVG]	Al ₂ O ₃ : 3.727; SiO ₂ : 44.12; CuO: 48.902; SO ₃ : 0.282; MnO: 0.247
P3 Plancheite	UP	Oberhä3	Jordan	slightly altered	P3 Plancheite UP-Oberhä3 [5x5 AVG]	Al ₂ O ₃ : 2.74; SiO ₂ : 43.25; CuO: 51.37; SO ₃ : 0.266; MnO: 0.085

Formatted: Font: 8 pt

Formatted Table

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 9 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 9 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 9 pt




Figure 10: Spectral library plot of the copper-bearing minerals – sulphides and native copper.




Figure 11: Spectral library plot of the copper-bearing minerals – silicates, carbonates and sulphates.

Formatted: Caption, Right: 0 cm, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Formatted: Caption, Right: 0 cm, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

4.3 Apliki mine samples

Table 12: Sample names, spectral library filenames, description and mineralogy of Apliki mine sample collection

Sample ID "Spectra name"	Description	Mineralogy based on qualitative XRD analysis (in no particular order) from Koerting (2021)
Apl1 A 1a, "Apl1 A 1a [5x5 AVGI]"	Grey-green "fresh" surface	Not available
Apl1 A 1b, "Apl1 A 1b [5x5 AVGI]"	Hematite coloured weathering crust	Andesine (anorthic), Quartz, Magnetite, Montmorillonite
Apl1 A 1d, "Apl1 A 1d [5x5 AVGI]"	"Fresh", dark green weathering crust	Anorthite, Magnetite, Diopside, Quartz, Montmorillonite
Apl1 A 1e, "Apl1 A 1e [5x5 AVGI]"	Yellow-ish orange weathering crust	Magnetite, Quartz, Montmorillonite, Diopside, Anorthite
Apl1 A 1f, "Apl1 A 1f [5x5 AVGI]"	"Soil formation", gravel	Magnetite, Anorthite, Quartz, Montmorillonite, Pyrite
Apl1 A 2a, "Apl1 A 2a [5x5 AVGI]"	Waste, soil	Goethite, Quartz, Clinocllore, Jarosite-Natrojarosite, Andesine, Gypsum
Apl1 A 3a, "Apl1 A 3a [5x5 AVGI]"	Yellow-ish weathered, soil	Andesine (anorthic), Quartz, Gypsum, Clinocllore, Jarosite, Montmorillonite
Apl1 A 3b, Apl1 A 3b [5x5 AVGI]	Brown-ish weathered, soil	Quartz, Andesine, Clinocllore, Gypsum, Jarosite, Montmorillonite
Apl1 A 4a, "Apl1 A 4a [5x5 AVGI]"	White, small grained gravel	Gypsum, Quartz, Clinocllore, Rozenite
Apl1 A 4b, "Apl1 A 4b [5x5 AVGI]"	Grey, small grained gravel	Quartz, Clinocllore, Andesine, Gypsum, Montmorillonite
Apl1 A 4c, "Apl1 A 4c [5x5 AVGI]"	Grey-green, weathering crust	Quartz, Clinocllore
Apl1 A 5a, "Apl1 A 5a [5x5 AVGI]"	Grey-medium, weathering crust	Gypsum, Quartz, Clinocllore
Apl1 A 5b, "Apl1 A 15b [5x5 AVGI]"	Grey-dark, weathering crust	Gypsum, Quartz, Clinocllore
Apl1 A 5c, "Apl1 A 5c [5x5 AVGI]"	Grey-light, weathering crust	Quartz, Gypsum, Clinocllore, Goethite, Hexahydrite
Apl1 A 6a, "Apl1 A 6a [5x5 AVGI]"	Red-ish brown, soil, gravel	Quartz, Pyrite, Analcime, Goethite, Montmorillonite, Clinocllore, Anorthite
Apl1 A 6b, "Apl1 A 6b [5x5 AVGI]"	Red-ish brown, soil	Anorthite, Quartz, Magnetite, Diopside, Montmorillonite, Gypsum, Goethite
Apl1 A 6c, "Apl1 A 6c [5x5 AVGI]"	Red-ish brown	Quartz, Clinocllore, Analcime, Gypsum, Calcite, Jarosite, Pyrite, Montmorillonite
Apl1 A 6d, "Apl1 A 6d [5x5 AVGI]"	Red-ish brown, soil	Quartz, Pyrite, Anorthite, Analcime, Clinocllore, Montmorillonite
Apl1 A 7d, "Apl1 A 7d [5x5 AVGI]"	Grey, crust unstable	Quartz, Hexahydrite, Clinocllore, Gypsum, Pyrite
Apl1 A 7d Hem, "Apl1 A 7d Hem [5x5 AVGI]"	Red, hematite	Pyrite, Hematite, Quartz, Gypsum, Clinocllore
Apl1 A 7e, "Apl1 A 7e [5x5 AVGI]"	Blue crystal	Rozenite, Goethite, Quartz, Apiohnite, Ferroxahydrite
Apl1 A 8a, "Apl1 A 8a [5x5 AVGI]"	Grey, small grained gravel	Quartz, Clinocllore, Pyrite, Ajoite
Apl1 A 8b, "Apl1 A 8b [5x5 AVGI]"	Grey, small grained gravel	Quartz, Clinocllore, Pyrite, Ajoite
Apl1 A 8c, "Apl1 A 8c [5x5 AVGI]"	Grey, soil-ish	Quartz, Clinocllore, Pyrite, Ajoite
Apl1 A 9a, "Apl1 A 9a [5x5 AVGI]"	Light green weathering crust	Quartz, Clinocllore (Mn), Clinocllore
Apl1 A 9b, "Apl1 A 9b [5x5 AVGI]"	Hematite vein	Quartz, Clinocllore, Pyrite, Hematite
Apl1 A 10a, "Apl1 A 10a [5x5 AVGI]"	White with pink, weathering crust	Clinocllore, Hematite, Quartz
Apl1 A 10b, "Apl1 A 10b [5x5 AVGI]"	White with purple, weathering crust	Quartz, Clinocllore
Apl1 A 10c, "Apl1 A 10c [5x5 AVGI]"	Green-ish veins	Quartz, Clinocllore

Formatted	... [90]
Formatted	... [91]
Formatted Table	... [92]
Formatted	... [93]
Formatted	... [94]
Formatted	... [95]
Formatted	... [96]
Formatted	... [97]
Formatted	... [98]
Formatted	... [99]
Formatted	... [100]
Formatted	... [101]
Formatted	... [102]
Formatted	... [103]
Formatted	... [104]
Formatted	... [106]
Formatted	... [107]
Formatted	... [105]
Formatted	... [108]
Formatted	... [109]
Formatted	... [110]
Formatted	... [111]
Formatted	... [112]
Formatted	... [113]
Formatted	... [114]
Formatted	... [115]
Formatted	... [116]
Formatted	... [117]
Formatted	... [118]
Formatted	... [119]
Formatted	... [121]
Formatted	... [122]
Formatted	... [123]
Formatted	... [120]
Formatted	... [124]
Formatted	... [125]
Formatted	... [126]
Formatted	... [127]
Formatted	... [128]
Formatted	... [129]
Formatted	... [130]
Formatted	... [131]
Formatted	... [132]
Formatted	... [133]
Formatted	... [134]
Formatted	... [135]
Formatted	... [136]
Formatted	... [137]
Formatted	... [138]
Formatted	... [139]
Formatted	... [140]
Formatted	... [141]
Formatted	... [142]
Formatted	... [143]

<u>ApI1 A 10d, "ApII A 10d [5x5 AVGI]"</u>	<u>White evaporitic crust</u>	<u>Quartz, Clinocllore, Pyrite</u>
<u>ApI1 A 11a, "ApII A 11a [5x5 AVGI]"</u>	<u>Grey, weathering crust</u>	<u>Quartz, Clinocllore, Gypsum, Bassanite</u>
<u>ApI1 A 11b, "ApII A 11b [5x5 AVGI]"</u>	<u>Green, weathering crust</u>	<u>Quartz, Clinocllore, Sphalerite</u>
<u>ApI1 A 13a, "ApII A 13a [5x5 AVGI]"</u>	<u>Red, rock</u>	<u>Andesine, Quartz, Magnetite, Montmorillonite-Chlorite, Diopside</u>
<u>ApI1 A 13b, "ApII A 13b [5x5 AVGI]"</u>	<u>Red, gravel, weathered hillside rock</u>	<u>Clinocllore, Quartz, Montmorillonite</u>
<u>ApI1 A 15a, "ApII A 15a [5x5 AVGI]"</u>	<u>Dark blue crystalline crust</u>	<u>Quartz (82.6%), Pyrite (7.5%), Chalcoppyrite (0.8%), Pentahydrate (cuprian) (9.1%)</u>
<u>ApI1 A 15b, "ApII A 15b [5x5 AVGI]"</u>	<u>Light blue rock and blue crust</u>	<u>Quartz (86.1%), Pyrite (4.5%), Pentahydrate (cuprian) (7.1%), Covellite (2.4%)</u>
<u>ApI1 A 15c, "ApII A 15c [5x5 AVGI]"</u>	<u>Black pyrite</u>	<u>Covellite (18.9%), Quartz (39.9%), Chalcanthite (21.8%), Pyrite (20.0%)</u>




Fig. 12: Spectral library of the 37 different Apliki mine samples, spectra stacked with offset.

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Not Bold

Formatted: Font: 8 pt

Formatted: Font: (Default) Calibri, 11 pt




Figure 13: Detailed, unstacked view of a selection of spectra, modified from (Koerting, 2021).

5. Validation and Discussion

Technical validation of the results in terms of sample material properties, systematic errors and variation of measurements (experimental error) are given below.

5.1 Sample Material Properties

The REO powders were certified to contain at least 99.9% of the corresponding REO. The certificates are listed in (Koerting et al., 2019a). The REE mineral samples were geochemically analysed using the Thermo Niton XL3t (Fisher Scientific, 2002) device. The resulting element concentrations and the measurement error (2σ) are provided in Koerting et al. (2019a). The validation for the copper-bearing minerals can be found in Koellner et al. (2019) and the Apliki mine sample validation, analyzed by BVM, can be found in Koerting et al. (2019b).

5.2 Systematic Errors of hyperspectral data acquisition

Systematic errors are discussed based on instrument drift, calibration and optimization of measurements. Initializing a warm-up phase of optical components, detectors and lamps, reduced influences due to instrument drift. Additionally, laboratory conditions were monitored to ensure a stable temperature and humidity. The HySpex cameras and the reference standards are factory calibrated once per year. Measurements used for the final reflectance spectral library were collected within one calibration time span to ensure equal acquisition conditions. For HySpex, averaging multiple measurements minimizes

Formatted: Font colour: Black

Formatted: Font: 9 pt, Bold

Formatted: Font: 9 pt, Bold

Formatted: Font: 9 pt, Bold, Not Italic

Formatted: Font: 9 pt, Bold, Not Italic

Deleted: 6

Deleted: z

variations in the data. An average (median) of 500 to 800 pixels [reflectance spectrum](#) was taken for the HySpex REE and REO [reflectance](#) spectra. This number relates to the maximum number of non-disturbed pixels per sample region of interest (e.g., pixels that were not shadowed from the sample holder side walls, etc.). For the copper-bearing minerals and the Apliki mine powders a 5x5 average pixel window was chosen over the area of interest. For these samples using a smaller pixel number for the average was necessary as the sampling of the copper-bearing minerals for geochemical validation occurred over a small area of the sample and the Apliki mine powder tablets were too small to ensure a larger homogenous area.

Deleted: a

5.3 Measurements variation

Variations of measurements were not only based on instrument calibrations or drift. They can also occur due to the detector geometry or geochemical properties of the minerals. These variations may appear as a shift of the peak positions of the absorption bands. This means, different hyperspectral sensors will show variations in the spectrum of the same material. By only using one set of hyperspectral sensors, the HySpex VNIR and SWIR, these shifts will not appear in our data sets. They might show when comparing our [reflectance](#) spectra of a material with [reflectance](#) spectra taken from a different instrument. For the copper-bearing minerals, the sample [reflectance](#) spectra also differ when comparing different samples of the same mineral species (e.g. “Malachite”) to each other. The spectral signal differs, for example, due to changes in geochemistry, physical appearance e.g. crystallization and degree of weathering (Clark, 1999; Hunt, 1982; Hunt and Ashley, 1979). [To avoid measurement variations caused by different sensors imaging data from the same sensors as the spectral library should to be used for the analysis. An example for an application can be using the here provided spectral library of the Apliki mine samples for an analysis of the HySpex hyperspectral imaging data of the Apliki mine face to be published in 2021 \(Koerting et al., 2021\).](#)

Deleted: 0

5.4 XL3t systematic errors

The XL3t is internally calibrated and provides an internal warm-up phase to guarantee stable measurement conditions. Unlike the spectrometer measurements, experimental error was only provided for the XL3t. In order to reduce the experimental error, a long duration measurement time of 120 seconds was set. The XL3t collects the emitted radiation from the sample using four different filters. While the sample was irradiated, each filter measures counts per second within a time span of 30 seconds. Next, the average counts per second were internally transformed to ppm. The irradiation of, in total, 120 seconds per sample was empirically tested to enable short measurement duration in combination with the lowest achievable standard deviation of concentration level.

6 Data Availability

The spectral libraries are published under the Creative Commons Attribution International 4.0 Licence (CC BY 4.0) via GFZ Data Services. Due to the different types of samples, we present the following three data publications: (1) [Mineral reflectance of 29 rare-earth minerals and rare-earth oxide powders including niobium- and tantalum-oxide powder, V. 2.0](#)

Deleted: 7

Deleted: 32

Deleted: (Koerting et al., 2019a)

660 GFZ Data Services, <http://doi.org/10.5880/GFZ.1.4.2019.004> (Koerting et al., 2019a); (2) Mineral reflectance spectra and chemistry of 20 copper-bearing minerals, V. 2.0 GFZ Data Services, <http://doi.org/10.5880/GFZ.1.4.2019.003> (Koellner et al., 2019), and (3) Mineral reflectance spectra and chemistry of 37 copper-bearing surface samples from Apliki copper-gold-pyrite mine in the Republic of Cyprus, V. 2.0 GFZ Data Services, <http://doi.org/10.5880/GFZ.1.4.2019.005> (Koerting et al., 2019b).

665 **7. Sample Availability**

The samples provided by the BGR are available through the collection of the BGR Spandau by their sample and collection name, in the technical report (<https://www.gewis.bgr.de>). The samples provided by the GFZ and UP belong to projects and have to be requested separately.

670 **8. Appendix**

Table A1: List of less commonly known terms and their abbreviations used throughout the paper

Terms	Abbreviation	Description
Abbreviation	REE	Rare Earth Element
	REO	Rare Earth Oxide
	REMin	Rare Earth Element-bearing Mineral
	VNIR	Visible light and near infrared
	SWIR	Short wave infrared
	XRF	X-Ray fluorescence
	EnMAP	Environmental Mapping and Analysis Program: future earth observation satellite mission (www.enmap.org) ¹
	CCRSS-A	China Commercial Remote Sensing Satellite System: future earth observation satellite mission
	HISUI	Hyperspectral Imager Suite: future earth observation satellite mission
Instruments	HySpex VNIR-1600	HySpex push broom spectrometer, VNIR camera
	HySpex SWIR-320m-e	HySpex push broom spectrometer, SWIR camera

Deleted:)
Deleted: Mineral
Deleted: (Koellner et al., 2019)
Deleted: ;
Formatted: Font: 10 pt
Deleted:
Deleted:)
Deleted: Mineral
Deleted: (Koerting et al., 2019b);
Deleted:)
Deleted: 8
Deleted: name
Deleted: table S5 in the supplements

Deleted: 9
Deleted: ces

Deleted:

	HySpex ground	HySpex operational software for laboratory and near-field application
	HySpex rad	HySpex calibration software to transform raw DN into radiance data
	Thermo Scientific Niton XL3t	Thermo Scientific Inc. X-Ray fluorescence analyzer (NITON TM XL3t)
	NDTr	Thermo Scientific Inc. NITON TM operational software
	JEOL JXA-8200	Electron probe microanalyzer (EPMA)
	JEOL JSM-6510	Scanning electron microscope (SEM)
	Oxford Instruments INCAx-act	Energy dispersive X-ray spectrometer (EDS)
Registered brands, Copyrights and/ or other protected terms	REacton®	Series of rare earth metals and compounds
	Alfa Aesar	Manufacturer and supplier of chemicals for research and development (today: "Thermo Scientific Inc.")
	Gunnar Färber Minerals	Supplier of mineral specimen
	REEMAP	Rare Earth Element MAPping: Research project for the development of a modular multi-sensor processing chain for modern imaging spectrometers to detect REEs
	Smithsonian Institution	Smithsonian Institution Department of Mineral Sciences, reference material from the Smithsonian Microbeam Standards
	Astimex Standards Ltd.	Astimex produces standards suitable for electron probe and scanning electron microscope X-ray analysis.
	BVM	Bureau Veritas Minerals is an industry leader in the analysis of minerals for the Exploration and Mining industries. BVM is a service-provider company that provides mineral preparation and laboratory testing services.
Research and	BGR	Federal Institute for Geosciences and Natural Resources

federal institutes	GSD	Geological Survey Department, Ministry of Agriculture, Rural Development and Environment, Republic of Cyprus
	UP	University of Potsdam
	GFZ	German Research Centre for Geosciences
Registered Trademarks	Excel™	Microsoft Excel™

690 **9. Author contributions**
Apliki mine and copper-bearing minerals: Friederike Koerting designed the Apliki sample related study, performed and ~~the~~ **measurements of the** Apliki samples and wrote the manuscript. Nicole Koellner designed the copper sample study, supervised the measurements and performed the geochemical analysis at the University of Potsdam. ~~Christian Mielke and~~ Agnieszka Kuras prepared parts of the spectral libraries. *REE minerals and REOs:* Nina K. Boesche designed the REE study, 695 performed some measurements, and supervised the REE measurements. Sabrina Herrmann prepared the samples and conducted most of the measurements. Christian Rogass developed and applied the HySpex post-processing chain. ~~Christian~~ Mielke and Kirsten Elger helped revising the manuscript. Uwe Altenberger supervised the studies and gave ~~valuable~~ comments on the manuscript.

700 **10. Competing interests**
The authors declare no conflict of interest

11. Acknowledgements
We would like to thank the Helmholtz Centre Potsdam GFZ German Research Centre for Geosciences for providing the 705 infrastructure and personnel support to conduct our research. Our gratitude also goes to the German Federal Ministry of Education and Research and the r4 subsidy program for innovative technologies for resource efficiency, which supported the REEMAP scientific project. We also thank the DLR Space Administration and the German Federal Ministry for Economic Affairs and Energy for the financial support based on a decision by the German Bundestag in the frame of the EnMAP scientific preparation program (Contract No. 50EE1256). We also want to express our gratitude to Seltenerdmetalle24, in 710 person Manuel Schultz, for his friendly service when providing laboratory standards and negative control sample holder. Thanks to the support by the GSD we were able to conduct a study and sample in the Republic of Cyprus and our thanks goes to our colleagues there for their help and directions in the unknown terrain. All the work in the Republic of Cyprus was conducted under the “Permit to conduct a Geological Survey, Ref. No. 02.13.005.002.005.022” from the 19th of March 2018, granted by the Geological Survey Department, Ministry of Agriculture, Rural Development and Environment (GSD) and the 715 Director Dr. Costas Constantinou. After the termination of the permit, a Memorandum of Understanding (MoU) and Framework for cooperation in the area of geo-science between the GSD and the GFZ was agreed upon in March 2019, the

- Deleted: 10
- Deleted: supervised the measurements for Marcel Hornings’ data and thethe measurements of the
- Moved down [4]: Marcel Horning performed most of the measurements on the copper samples and prepared the spectral copper library. Pia Brinkm
- Moved down [3]: Pia Brinkman prepared the Apliki sample powder tablets. Christian Mielke
- Deleted: ,
- Deleted: , Constantin Hildebrand and Friederike Klos
- Deleted:
- Deleted: ¶
- Deleted: some
- Deleted: 1
- Deleted: 2

publication of the Apliki mine surface data is associated to this MoU. [Constantin Hildebrand and Friederike Klos prepared parts of the spectral libraries for the data publications and contributed insight into the spectral interpretation.](#) [Pia Brinkman prepared the Apliki sample powder tablets.](#) [Marcel Horning performed most of the measurements on the copper samples and prepared the spectral copper library during his B.Sc. thesis.](#) We thank our colleagues for their input and insights.

12 References

Baldrige, A. M., Hook, S. J., Grove, C. I. and Rivera, G.: The ASTER spectral library version 2.0, Remote Sensing of Environment, 113(4), 711–715, doi:10.1016/j.rse.2008.11.007, 2009.

Boesche, N., Rogass, C., Lubitz, C., Brell, M., Hermann, S., Mielke, C., Tonn, S., Appelt, O., Altenberger, U. and Kaufmann, H.: Hyperspectral REE (Rare Earth Element) Mapping of Outcrops—Applications for Neodymium Detection, Remote Sensing, 7(5), 5160–5186, doi:10.3390/rs70505160, 2015.

Boesche, N. K., Rogass, C., Mielke, C., Hermann, S., Körting, F., Papenfuß, A., Lubitz, C., Brell, M., Tonn, S. and Altenberger, U.: Chapter 16 - Hyperspectral Rare Earth Element Mapping of Three Outcrops at the Fen Complex, Norway: Calcitic, Dolomitic, and Ankeritic Carbonatites, in Rare Earths Industry - Technological, Economic, and Environmental Implications, p. 235-265, Elsevier Inc., <https://doi.org/10.1016/B978-0-12-802328-0.00016-4>, 2016.

Bösche, N. K.: Detection of Rare Earth Elements and Rare Earth Oxides with Hyperspectral Spectroscopy, University of Potsdam. [online] Available from: <http://nbn-resolving.de/urn:nbn:de:kobv:517-opus4-85363>, 2015.

Bureau Veritas: Bureau Veritas Metals, Minerals & Environmental Schedule of Services & Fees 2020. [online] Available from: http://acmelab.com/wp-content/uploads/2020/02/BV_Fees-Schedule-2020_USD_v3_07Feb2020.pdf, 2020.

Clark, R. N.: Spectroscopy of rocks and minerals, and principles of spectroscopy, Remote sensing for the earth sciences: Manual of remote sensing, 3, 3–58, doi:10.1111/j.1945-5100.2004.tb00079.x, 1999.

Clark, R. N., Swayze, G. A., Wise, R. A., Livo, K. E., Hoefen, T. M., Kokaly, R. F. and Sutley, S. J.: USGS Spectral Library splib06a: U.S. Geological Survey, Digital Data Series 231, <http://speclab.cr.usgs.gov/spectral.lib06>, [online] Available from: <http://speclab.cr.usgs.gov/spectral.lib06>, 2007.

Guanter, L., Kaufmann, H., Segl, K., Foerster, S., Rogass, C., Chabrillat, S., Kuester, T., Hollstein, A., Rossner, G., Chlebek, C., Straif, C., Fischer, S., Schrader, S., Storch, T., Heiden, U., Mueller, A., Bachmann, M., Mühle, H., Müller, R., Habermeyer, M., Ohndorf, A., Hill, J., Buddenbaum, H., Hostert, P., Van Der Linden, S., Leitão, P. J., Rabe, A., Doerffer, R., Krasemann, H., Xi, H., Mauser, W., Hank, T., Locherer, M., Rast, M., Staenz, K. and Sang, B.: The EnMAP spaceborne imaging spectroscopy mission for earth observation, Remote Sensing, doi:10.3390/rs70708830, 2015.

Heinrich, K. F. J. and Newbury, D. E.: Electron probe quantitation edited by K.F.J. HEINRICH and D.E. NEWBURY (Plenum Press New York, 1991), Microscopy Microanalysis Microstructures, doi:10.1051/mm:0199200302-3029500, 1991.

Herrmann, S.: Capacity of Imaging Spectroscopy for the characterisation of REO , REE bearing minerals & primary REE-deposits., Master Thesis, (Scientific Technical Report ; 19/08), Potsdam : GFZ German Research Centre for Geosciences, vii, 161p., <https://doi.org/10.2312/GFZ.b1v03-19089>, 2019.

Hunt, G. R.: Spectral signatures of particulate minerals in the visible and near infrared, Geophysics, doi:10.1190/1.1440721, 1977.

Moved (insertion) [3]

Moved (insertion) [4]

Deleted: ¶

¶

Deleted: 3

Formatted: Font: 9 pt, Not Bold, Font colour: Auto

Formatted: Indent: Left: 0 cm, First line: 0 cm, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Deleted: edited by I. B. De Lima and W. L. Filho,

Deleted: 434

Deleted: [online] Available from: <https://www.elsevier.com/books/rare-earths-industry/borges-de-lima/978-0-12-802328-0>, 2015.¶

- Hunt, G. R.: Spectroscopic Properties of Rocks and Minerals, in Practical Handbook of Physical Properties of Rocks and Minerals, edited by R. S. Carmichael, pp. 599–669, CRC Press., 1982.
- Hunt, G. R. and Ashley, R. P.: Spectra of altered rocks in the visible and near infrared, *Economic Geology*, doi:10.2113/gsecongeo.74.7.1613, 1979.
- 780 hypspec.no/products/disc.php: Norsk Elektro Optikk AS HySpex VNIR1600 and SWIR320 m-e, [online] Available from: <https://www.hypspec.no/products/disc/vnir-1600.php> (Accessed 18 June 2019), 2019.
- Iwasaki, A., Ohgi, N., Tanii, J., Kawashima, T. and Inada, H.: Hyperspectral Imager Suite (HISUI)-Japanese hyper-multi spectral radiometer, in *International Geoscience and Remote Sensing Symposium (IGARSS)*., 2011.
- Jensen, J. R.: *Remote Sensing of the Environment An Earth Resource Perspective*, 2nd ed., Pearson Education., 2010.
- 785 Koellner, N., Koerting, F., Horning, M., Mielke, C. and Altenberger, U.: [Technical Report: Mineral reflectance spectra](#) and chemistry of 20 copper-bearing minerals., [V. 2.0 GFZ Data Services](#), <https://doi.org/10.5880/GFZ.1.4.2019.003>, 2019.
- Koerting, F., Herrmann, S., Boesche, N. K., Mielke, C., Koellner, N. and Altenberger, U.: [Technical Report: Mineral reflectance spectra](#) and chemistry of [29 rare-earth minerals and rare-earth oxide powders including niobium- and tantalum-oxid powder.](#), [V. 2.0 GFZ Data Services](#), <https://doi.org/10.5880/GFZ.1.4.2019.004>, 2019a.
- 790 Koerting, F., Rogass, C., Koellner, N., Horning, M. and Altenberger, U.: [Technical Report: Mineral reflectance spectra](#) and chemistry of 37 copper-bearing surface samples from Apliki copper-gold-pyrite mine in the Republic of Cyprus., [V. 2.0 GFZ Data Services](#), <https://doi.org/10.5880/GFZ.1.4.2019.005>, 2019b.
- Koerting, F., Koellner, N., Mielke, C., Rogass, C., Kuras, A., Altenberger, U., Klos, F. and Hildebrand, C.: Hyperspectral mine face scan of the northern open cut and laboratory scan of surface from the Apliki copper-gold-pyrite mine in the Republic of Cyprus, [2021](#).
- 795 Koerting, F. M.: Hybrid imaging spectroscopy approaches for open pit mining - Applications for virtual mine face geology, [Doctoral Thesis](#), University of Potsdam, [2021](#).
- Kokaly, R. F., Clark, R. N., Swayze, G. A., Livo, K. E., Hoefen, T. M., Pearson, N. C., Wise, R. A., Benzel, W. M., Lowers, H. A., Driscoll, R. L. and Klein, A. J.: USGS Spectral Library Version 7. [online] Available from: <https://pubs.er.usgs.gov/publication/ds1035>, 2017.
- 800 Körting, F.: Development of a 360 ° hyperspectral drill core scanner Test of technical conditions and validation of high-resolution near-field analysis of crystalline basement rocks using COSC-1 core samples., [Master Thesis. \(Scientific Technical Report : 19/07\). Potsdam : GFZ German Research Centre for Geosciences, 97, xxi p., https://doi.org/10.2312/GFZ.b103-19071](#), 2019.
- Lorenz, M., Altenberger, U., Trumbull, R. B., Lira, R., Luchi, M. L. De, Günter, C. and Eidner, S.: Chemical and textural relations of britholite- And apatite-group minerals from hydrothermal REE mineralization at the Rodeo de los Molles deposit, Central Argentina, *American Mineralogist*, doi:10.2138/am-2019-6969, 2019.
- 805 van der Meer, F. D., van der Werff, H. M. A., van Ruitenbeek, F. J. A., Hecker, C. A., Bakker, W. H., Noomen, M. F., van der Meijde, M., Carranza, E. J. M., de Smeth, J. B. and Woldai, T.: Multi- and hyperspectral geologic remote sensing: A review, *International Journal of Applied Earth Observation and Geoinformation*, doi:10.1016/j.jag.2011.08.002, 2012.
- Meerdink, S. K., Hook, S. J., Roberts, D. A. and Abbott, E. A.: The ECOSTRESS spectral library version 1.0, *Remote Sensing of Environment*, doi:10.1016/j.rse.2019.05.015, 2019.
- 810 Mielke, C., Rogass, C., Boesche, N., Segl, K. and Altenberger, U.: EnGeoMAP 2.0—Automated Hyperspectral Mineral Identification for the German EnMAP Space Mission, *Remote Sensing*, 8(2), 127, doi:10.3390/rs8020127, 2016.
- Percival, J. B., Olejarz, A. D., English, M. L. R., Belley, P. M., Flynn, T., Laudadio, A. B., Stirling, J. A. and R: *The National Mineral*

Deleted: Mineral spectra

Deleted: Mineral spectra

Deleted: 32

Deleted: 2020.

Deleted: 2021

- Reference Collection (NMC) Digital Spectral (VIS-NIR-SWIR) Library. Part I: The Kodama clay mineral collection., 2016.
- 820 Rogass, C., Koerting, F. M., Mielke, C., Brell, M., Boesche, N. K., Bade, M. and Hohmann, C.: Translational imaging spectroscopy for proximal sensing, *Sensors (Switzerland)*, 17(8), doi:10.3390/s17081857, 2017.
- Swayze, G. A., Clark, R. N., Goetz, A. F. H., Livo, K. E., Breit, G. N., Kruse, F. A., Sutley, S. J., Snee, L. W., Lowers, H. A., Post, J. L., Stoffregen, R. E. and Ashley, R. P.: Mapping advanced argillic alteration at Cuprite, Nevada, using imaging spectroscopy, *Economic Geology*, 109(5), 1179–1221, doi:10.2113/econgeo.109.5.1179, 2014.
- 825 Tong, Q., Xue, Y. and Zhang, L.: Progress in hyperspectral remote sensing science and technology in China over the past three decades, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, doi:10.1109/JSTARS.2013.2267204, 2014.
- Turner, D., Rivard, B. and Groat, L.: Rare earth element ore grade estimation of mineralized drill core from hyperspectral imaging spectroscopy, in 2014 IEEE Geoscience and Remote Sensing Symposium, pp. 4612–4615, IEEE., 2014a.
- Turner, D. J.: Reflectance spectroscopy and imaging spectroscopy of rare earth element-bearing mineral and rock samples., 2015.
- 830 Turner, D. J., Rivard, B. and Groat, L. A.: Visible and short-wave infrared reflectance spectroscopy of REE fluorocarbonates, *American Mineralogist*, doi:10.2138/am.2014.4674, 2014b.

Deleted:

Boesche, N., Rogass, C., Lubitz, C., Brell, M., Herrmann, S., Mielke, C., Tonn, S., Appelt, O., Altenberger, U. and Kaufmann, H.: Hyperspectral REE (Rare Earth Element) Mapping of Outcrops—Applications for Neodymium Detection, *Remote Sensing*, 7(5), 5160–5186, doi:10.3390/rs70505160, 2015.

Boesche, N. K.: Hyperspectral Rare Earth Element Mapping of Three Outcrops at the Fen Complex, Norway: Calcitic, Dolomitic, and Ankeritic Carbonatites, in *Rare Earths Industry - Technological, Economic, and Environmental Implications*, edited by I. B. De Lima and W. L. Filho, p. 434, Elsevier Inc. [online] Available from: <https://www.elsevier.com/books/rare-earths-industry/borges-de-lima/978-0-12-802328-0>, 2015.

Bösch, N. K.: Detection of Rare Earth Elements and Rare Earth Oxides with Hyperspectral Spectroscopy, University of Potsdam. [online] Available from: <http://nbn-resolving.de/urn:nbn:de:kobv:517-opus4-85363>, 2015.

Bureau Veritas: Bureau Veritas Metals, Minerals & Environmental Schedule of Services & Fees 2020. [online] Available from: http://acmelab.com/wp-content/uploads/2020/02/BV_Fees-Schedule-2020_USD_v3_07Feb2020.pdf, 2020.

Clark, R. N.: Spectroscopy of rocks and minerals, and principles of spectroscopy, *Remote sensing for the earth sciences: Manual of remote sensing*, 3, 3–58, doi:10.1111/j.1945-5100.2004.tb00079.x, 1999.

Clark, R. N., Swayze, G. A., Wise, R. A., Livo, K. E., Hoefen, T. M., Kokaly, R. F. and Sutley, S. J.: USGS Spectral Library splib06a: U.S. Geological Survey, Digital Data Series 231, <http://speculab.cr.usgs.gov/spectral.lib06>, [online] Available from: <http://speculab.cr.usgs.gov/spectral.lib06>, 2007.

Guanter, L., Kaufmann, H., Segl, K., Foerster, S., Rogass, C., Chabrilat, S., Kuester, T., Hollstein, A., Rossner, G., Chlebek, C., Straif, C., Fischer, S., Schrader, S., Storch, T., Heiden, U., Mueller, A., Bachmann, M., Mühle, H., Müller, R., Habermeyer, M., Ohndorf, A., Hill, J., Buddenbaum, H., Hostert, P., Van Der Linden, S., Leitão, P. J., Rabe, A., Doerffer, R., Krasemann, H., Xi, H., Mauser, W., Hank, T., Locherer, M., Rast, M., Staenz, K. and Sang, B.: The EnMAP spaceborne imaging spectroscopy mission for earth observation, *Remote Sensing*, doi:10.3390/rs70708830, 2015.

Heinrich, K. F. J. and Newbury, D. E.: *Electron probe quantitation* edited by K.F.J. HEINRICH and D.E. NEWBURY (Plenum Press New York, 1991), *Microscopy Microanalysis Microstructures*, doi:10.1051/mm:0199200302-3029500, 1991.

Herrmann, S.: Capacity of Imaging Spectroscopy for the characterisation of REO, REE bearing minerals & primary REE-deposits., 2019.

Hunt, G. R.: Spectral signatures of particulate minerals in the visible and near infrared, *Geophysics*, doi:10.1190/1.1440721, 1977.

Hunt, G. R.: *Spectroscopic Properties of Rocks and Minerals*, in *Practical Handbook of Physical Properties of Rocks and Minerals*, edited by R. S. Carmichael, pp. 599–669, CRC Press., 1982.

Hunt, G. R. and Ashley, R. P.: Spectra of altered rocks in the visible and near infrared, *Economic Geology*, doi:10.2113/gsecongeo.74.7.1613, 1979.

[hypex.no/products/disc.php](https://www.hypex.no/products/disc.php): Norsk Elektro Optikk AS HySpex VNIR1600 and SWIR320 m-e., [online] Available from: <https://www.hypex.no/products/disc/vnir-1600.php> (Accessed 18 June 2019), 2019.

Iwasaki, A., Ohgi, N., Tanii, J., Kawashima, T. and Inada, H.: Hyperspectral Imager Suite (HISUI)-Japanese hyper-multi spectral radiometer, in *International Geoscience and Remote Sensing Symposium (IGARSS)*, 2011.

Jensen, J. R.: Remote Sensing of the Environment An Earth... [174]

Page 10: [1] Deleted	Friederike Körting	27/10/2020 19:38:00
Page 10: [2] Deleted	Friederike Körting	27/10/2020 19:38:00
Page 10: [3] Deleted	Friederike Körting	27/10/2020 19:37:00
Page 10: [4] Deleted	Friederike Körting	27/10/2020 19:39:00
Page 10: [5] Deleted	Friederike Körting	27/10/2020 19:39:00
Page 10: [6] Deleted	Friederike Körting	27/10/2020 19:37:00
Page 10: [7] Deleted	Friederike Körting	27/10/2020 19:39:00
Page 10: [8] Deleted	Friederike Körting	27/10/2020 19:39:00
Page 10: [9] Deleted	Friederike Körting	27/10/2020 19:37:00
Page 13: [10] Deleted	Friederike Körting	27/10/2020 19:54:00
Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00

Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00
-----------------------	--------------------	---------------------

Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00
-----------------------	--------------------	---------------------

Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00
-----------------------	--------------------	---------------------

Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00
-----------------------	--------------------	---------------------

Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00
-----------------------	--------------------	---------------------

Page 14: [11] Deleted	Friederike Körting	19/12/2020 14:38:00
-----------------------	--------------------	---------------------

Page 14: [12] Formatted	Friederike Körting	19/12/2020 15:00:00
-------------------------	--------------------	---------------------

Heading 3

Page 14: [13] Formatted	Friederike Körting	26/10/2020 12:35:00
-------------------------	--------------------	---------------------

Font colour: Auto

Page 14: [14] Formatted	Friederike Körting	26/10/2020 13:14:00
-------------------------	--------------------	---------------------

Caption

Page 14: [15] Formatted	Friederike Körting	17/12/2020 23:36:00
-------------------------	--------------------	---------------------

Font: 8 pt

▲
Page 14: [18] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [19] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [20] Formatted Friederike Körting 17/12/2020 23:54:00

Font: 8 pt, German

▲
Page 14: [21] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [22] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [23] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [24] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [25] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [26] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [27] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [28] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [29] Formatted Table Friederike Körting 19/12/2020 14:44:00

Formatted Table

▲
Page 14: [30] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [31] Formatted Friederike Körting 17/12/2020 23:36:00

Font: 8 pt

▲
Page 14: [32] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [33] Formatted Friederike Körting 17/12/2020 23:39:00

Left

▲
Page 14: [36] Formatted **Friederike Körting** **17/12/2020 23:36:00**

Font: 8 pt

▲
Page 14: [37] Formatted **Friederike Körting** **17/12/2020 23:39:00**

Left

▲
Page 14: [38] Formatted **Friederike Körting** **17/12/2020 23:36:00**

Font: 8 pt

▲
Page 14: [39] Formatted Table **Friederike Körting** **19/12/2020 14:44:00**

Formatted Table

▲
Page 14: [40] Formatted **Friederike Körting** **17/12/2020 23:39:00**

Left

▲
Page 14: [41] Formatted **Friederike Körting** **17/12/2020 23:36:00**

Font: 8 pt

▲
Page 14: [42] Formatted **Friederike Körting** **17/12/2020 23:39:00**

Left

▲
Page 14: [43] Formatted **Friederike Körting** **17/12/2020 23:36:00**

Font: 8 pt

▲
Page 14: [44] Formatted **Friederike Körting** **17/12/2020 23:39:00**

Left

▲
Page 14: [45] Formatted **Friederike Körting** **17/12/2020 23:36:00**

Font: 8 pt

▲
Page 14: [46] Formatted **Friederike Körting** **17/12/2020 23:39:00**

Left

▲
Page 14: [47] Formatted **Friederike Körting** **17/12/2020 23:36:00**

Font: 8 pt

▲
Page 14: [48] Formatted Table **Friederike Körting** **19/12/2020 14:45:00**

Formatted Table

▲
Page 14: [49] Formatted **Friederike Körting** **17/12/2020 23:39:00**

Left

▲
Page 15: [50] Formatted **Friederike Körting** **17/12/2020 23:33:00**

Caption

▲
Page 15: [51] Formatted **Friederike Körting** **17/12/2020 23:35:00**

Font: 8 pt

Page 15: [54] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [55] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [56] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [57] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [58] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [59] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [60] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [61] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [62] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [63] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [64] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [65] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [66] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [67] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [68] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [69] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

Page 15: [72] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [73] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [74] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [75] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [76] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [77] Formatted	Friederike Körting	17/12/2020 23:54:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt, German

▲

Page 15: [78] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [79] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [80] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [81] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [82] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [83] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [84] Formatted	Friederike Körting	17/12/2020 23:54:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt, German

▲

Page 15: [85] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 15: [86] Formatted	Friederike Körting	17/12/2020 23:39:00
--------------------------------	---------------------------	----------------------------

Left

▲

Page 15: [87] Formatted	Friederike Körting	17/12/2020 23:35:00
--------------------------------	---------------------------	----------------------------

Font: 8 pt

Caption, Right: 0 cm, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border),
Between : (No border)

▲
Page 19: [90] Formatted **Friederike Körting** **26/10/2020 13:15:00**

Caption

▲
Page 19: [91] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [92] Formatted Table **Friederike Körting** **19/12/2020 14:51:00**

Formatted Table

▲
Page 19: [93] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [93] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [94] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [94] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [95] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [96] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [96] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [97] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [97] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [98] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [99] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [99] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

Page 19: [102] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt, Italic

▲
Page 19: [103] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [104] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [104] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [105] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [106] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt, Italic

▲
Page 19: [107] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [108] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [108] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [109] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [109] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [110] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [111] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [111] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [112] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [113] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

Page 19: [114] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [115] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [116] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [116] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [117] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [117] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [118] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [119] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [120] Formatted **Friederike Körting** **17/12/2020 15:18:00**

Tab stops: Not at 3,42 cm

▲
Page 19: [121] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt, Not Bold

▲
Page 19: [122] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [122] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [123] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [124] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [124] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [125] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

Page 19: [127] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [127] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [128] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [128] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [129] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [130] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [130] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [131] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [131] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [132] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [133] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [133] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [134] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [134] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [135] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [136] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

Page 19: [137] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [138] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [139] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [139] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [140] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [141] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [142] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [143] Formatted **Friederike Körting** **19/12/2020 14:52:00**

Indent: First line: 0 cm

▲
Page 19: [144] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [145] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [145] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [146] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [147] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [148] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [148] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

▲
Page 19: [149] Formatted **Friederike Körting** **17/12/2020 16:05:00**

Font: 8 pt

Page 19: [151] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [151] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [152] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [153] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [153] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [154] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [154] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [155] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [156] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [156] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [157] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [157] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [158] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [159] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [159] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

▲

Page 19: [160] Formatted	Friederike Körting	17/12/2020 16:05:00
---------------------------------	---------------------------	----------------------------

Font: 8 pt

Page 19: [162] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [162] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [163] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [163] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [164] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [165] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [165] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [166] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [166] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [167] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [168] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [168] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [169] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [170] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [170] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

▲

Page 19: [171] Formatted	Friederike Körting	17/12/2020 16:05:00
--------------------------	--------------------	---------------------

Font: 8 pt

Font: 8 pt